


Chapter 2.5 - Continuous Functions

Motivation for Continuous Function

Recall: For polynom $P(x)$ holds $\lim_{x \rightarrow a} P(x) = P(a)$.

What should happen at $x \rightarrow a$ happens at $x = a$.



This is a nice property and behavior, called *continuous*.

$f(x)$ is *continuous* at $x = a$ if $\lim_{x \rightarrow a} f(x) = f(a)$

Some Continuous Functions

If f is continuous at a , the following three MUST ALL happen:

- ▶ $\lim_{x \rightarrow a} f(x)$ exists
- ▶ $f(a)$ exists
- ▶ $\lim_{x \rightarrow a} f(x) = f(a)$

If any of them fail, f is *not* continuous at a .

The following are continuous at all points

- ▶ Polynomials
- ▶ Rational functions where denominator $\neq 0$.
- ▶ $\sin x$, $\cos x$, e^x ,
- ▶ $\ln x$ for $x > 0$

Intuition: If you can draw the graph of f around a with one stroke, then f is continuous at a .

Types of Discontinuity

Removable by (re)defining the function at $x = a$, we can fix the problem.

Example: Is $f(x) = (x^2 - 1)/(x - 1)$ continuous at $x = 1$?

It is not defined at $x = 1$, but

$$\lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 1} = 2.$$

Removable discontinuities are great because we can “fix” them. The piecewise-defined function

$$F(x) = \begin{cases} \frac{x^2 - 1}{x - 1} & x \neq 1 \\ 2 & x = 1 \end{cases}$$

is basically the function f , but with the open hole at $x = 1$ filled in.

Jump, left and right hand limits exist but disagree.

Example:

$$f(x) = \begin{cases} x + 3 & x < 1 \\ 2 & x = 1 \\ 2x - 2 & x > 1 \end{cases}$$

at $x = 1$.

$$\lim_{x \rightarrow 1^-} f(x) = 4$$

$$\lim_{x \rightarrow 1^+} f(x) = 0$$

Hence $\lim_{x \rightarrow 1} f(x)$ does not exist.

More Types of Discontinuity

Oscilating if limit does not exist due to oscilating.

Example:

$$f(x) = \begin{cases} \sin\left(\frac{1}{x}\right) & x > 0 \\ 0 & x \leq 0 \end{cases}$$

at $x = 0$.

$$\lim_{x \rightarrow 0^-} f(x) = 0$$

$$\lim_{x \rightarrow 0^+} f(x) \text{ does not exist}$$

Hence $\lim_{x \rightarrow 0} f(x)$ does not exist.

Infinite Discontinuity appears when asymptote from one (or both) sides. More in next section.

Example:

$$f(x) = \begin{cases} \frac{1}{x^2} & x \neq 0 \\ 0 & x = 0 \end{cases}$$

at $x = 0$.

$$\lim_{x \rightarrow 0^-} f(x) = \infty$$

$$\lim_{x \rightarrow 0^+} f(x) = \infty$$

Hence one could say $\lim_{x \rightarrow 0} f(x) = \infty$ but we cannot have $f(x) = \infty$.

Continuous Function

A function is *continuous* if it is continuous at *every* point.

Intuition: If you could draw the graph of f with one stroke.

Example: Find a and b so that $f(x)$ is continuous and

$$f(x) = \begin{cases} & x < -1 \\ ax + b & -1 \leq x \leq 1 \\ & x > 1 \end{cases}$$

Combining Continuous Functions

If $f(x)$ and $g(x)$ are continuous then so are

- ▶ $f(x) + g(x)$
- ▶ $Kf(x)$ for $K \in \mathbb{R}$
- ▶ $f(x)g(x)$
- ▶ $\frac{f(x)}{g(x)}$ if $g(x) \neq 0$
- ▶ $(f(x))^q$ for reasonable q
- ▶ $f(g(x))$ composition

Example: Is $f(x)$ continuous?

$$f(x) = \frac{\sin(e^x + \pi - 1) + x^3 - 16 \cos(\sin(x))}{x^4 + 2 + \sin^2(e^x - e^{-x}) + \ln(x^2 + 1)}$$

Yes. The functions $\sin x$, e^x , $\cos x$ are continuous. Also $\ln x$ is continuous for $x > 0$ and notice that $x^2 + 1 \geq 1$. In particular $\ln x^2 + 1 \geq 0$. Hence the denominator is > 0 and $f(x)$ is continuous.

Nice fact about continuous functions:


$$\lim_{x \rightarrow a} f(x) = f(a)$$

Example: Given $f(x)$ above find

$$\lim_{x \rightarrow 0} f(x) = \frac{\sin(\pi) + 0 - 16 \cos(0)}{0 + 2 + \sin^2(0) + \ln(1)} = -8$$

Intermediate Value Theorem

If $f(x)$ is *continuous* on $[a, b]$, then for any y between $f(a)$ and $f(b)$ there is a c in $[a, b]$ with $f(c) = y$.



Example: $f(x) = 18x^3 - 63x^2 + 67x - 20$
Does $f(x)$ have a root for $0 \leq x \leq 1$? $f(x)$ is continuous. Also $f(0) = -20$ and $f(1) = 2$. The IVT implies there is $c \in [0, 1]$ such that $f(c) = 0$. Actually, $c = \frac{1}{2}$.

Does $f(x)$ have a root for $1 \leq x \leq 2$? $f(1) = 2$ and $f(2) = 6$. The IVT does not say anything about the roots! There are two roots $\frac{4}{3}$ and $\frac{5}{3}$.

Example: Show exists $0 \leq x$ that $\cos(x) = x$. Let $f(x) = \cos(x) - x$. Using $x = 0$ and $x = \frac{\pi}{2}$ yields

$$f(0) = \cos(0) - 0 = 1 \quad f\left(\frac{\pi}{2}\right) = \cos\left(\frac{\pi}{2}\right) - \frac{\pi}{2} = -\frac{\pi}{2}$$


As 0 in $[-\frac{\pi}{2}, 1]$, the intermediate value theorem implies that there exists a c in $[0, \frac{\pi}{2}]$ such that $0 = f(c) = \cos(c) - c$ and thus $\cos(c) = c$.

$$\cos(\cos(\cos(\dots \cos(x) \dots)))$$

What happens when you type to calculator 0 and keep pressing \cos button? (in Rad)

Let's move between points

$(0, 0)$, $(0, \cos(0))$, $(\cos(0), \cos(0))$, $(\cos(0), \cos(\cos(0)))$, $(\cos(\cos(0)), \cos(\cos(0)))$, \dots



Notice the green spiral is behaving like a fractal.

Chapter 2.5 Recap

- ▶ $f(x)$ is *continuous at $x = a$* if $\lim_{x \rightarrow a} f(x) = f(a)$
- ▶ f is continuous if it is continuous at every point
- ▶ Removable discontinuity, jump, oscillation, asymptotes (next time)
- ▶ combining continuous functions gives continuous function
- ▶ Intermediate Value Theorem