

## Math-484 Homework #2

I will finish the homework before 11 am Sep 9 and bring it to class. If I have troubles with my work I may come to office hours or ask by email. If I spot a mathematical mistake I will let the lecturer know as soon as possible.

I will write clearly and neatly as the grader is not an expert in cryptography. I will sign each paper of my work and indicate if I am D14 (4 hours student).

**Exercise 1:** (What is positive/negative (semi)definite?)

Decide if the following matrices are positive or negative (semi)definite or indefinite and explain why:

$$(a) \begin{pmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 5 \end{pmatrix} \quad (b) \begin{pmatrix} 3 & 1 & 2 \\ 1 & 5 & 3 \\ 2 & 3 & 7 \end{pmatrix}$$
$$(c) \begin{pmatrix} -4 & 0 & 1 \\ 0 & -3 & 2 \\ 1 & 2 & -5 \end{pmatrix} \quad (d) \begin{pmatrix} 2 & 2 & 0 \\ 2 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

---

**Exercise 2:** (I will recall what is a quadratic form.)

Write the quadratic form  $Q_A(\mathbf{x})$  associated to matrix

$$A = \begin{pmatrix} -4 & 0 & 1 \\ 0 & -3 & 2 \\ 1 & 2 & -5 \end{pmatrix}.$$

---

**Exercise 3:** (I will recall what is coercive.)

Decide which of these functions  $\mathbb{R}^3 \rightarrow \mathbb{R}$  are coercive (of course, argue why):

$$(a) f(x, y, z) = x^3 + y^3 + z^3 - xy \quad (b) f(x, y, z) = x^4 + y^4 + z^2 - 3xy - z$$
$$(c) f(x, y, z) = x^4 + y^4 + z^2 - xyz^2 \quad (d) f(x, y, z) = x^4 + y^4 - 2xy^2$$

---

**Exercise 4:** (Do I understand the assumption of Theorem 8?)

Show that the principal minors of the matrix

$$A = \begin{pmatrix} 1 & -8 \\ 1 & 1 \end{pmatrix}$$

are positive, but there are  $\mathbf{x} \neq \mathbf{0}$  in  $\mathbb{R}^2$  such that  $\mathbf{x} \cdot A\mathbf{x} < 0$ . Why does this not contradict Theorem 8 (1.3.3 in the textbook)?

---

**Exercise 5:** (Can I use all that stuff to find minimizers and maximizers?)

Find (local, global) minimizers and maximizers of the following functions:

$$(a) f(x_1, x_2) = e^{-(x_1^2 + x_2^2)} \quad (b) f(x_1, x_2, x_3) = (2x_1 - x_2)^2 + (x_2 - x_3)^2 + (x_3 - 1)^2$$

---

**Exercise 6:** (A bit more coercive thinking. **D14 only**)

Find a function  $f(x, y)$  on  $\mathbb{R}^2$  such that for each real number  $t$ , we have

$$\lim_{x \rightarrow \infty} f(x, tx) = \lim_{y \rightarrow \infty} f(ty, y) = +\infty$$

but such that  $f(x, y)$  is not coercive.

---